



Gibbs versus non-Gibbs distributions in money dynamics

Marco Patriarca^{a,*}, Anirban Chakraborti^b, Kimmo Kaski^a

^aComplex Systems Group, Laboratory of Computational Engineering,
Helsinki University of Technology, P.O. Box 9203, HUT 02015, Finland

^bDepartment of Physics, Brookhaven National Laboratory, Upton, NY 11973, USA

Abstract

We review a simple model of closed economy, where the economic agents make money transactions and a saving criterion is present. We observe the Gibbs distribution for zero saving propensity, and non-Gibbs distributions otherwise. While the exact solution in the case of zero saving propensity is already known to be given by the Gibbs distribution, here we provide the explicit analytical form of the equilibrium distribution for the general case of nonzero saving propensity. We verify it through comparison with numerical data and show that it can be cast in the form of a gamma-distribution.

© 2004 Elsevier B.V. All rights reserved.

PACS: 89.65.Gh; 87.23.Ge; 02.50.-r

Keywords: Econophysics; Money dynamics; Gamma distribution; Gibbs distribution

1. Introduction

It is known that the higher end of the distribution of income $f(m)$ follows the Pareto law [1], $f(m) \propto m^{-1-\alpha}$, where m is the income (money) and the exponent α has a value between 1 and 2 [2–5]. An explanation of the Pareto law, in terms of the laws regulating the system micro-dynamics, should take into account its basic constituents, i.e., the trading agents, as well as the criteria used to carry out the economic transactions. Several studies have been made to provide an explanation (see Ref. [6] for a brief summary and more references). In this respect, it is of general interest to study

* Corresponding author. Fax: +358-9-451-4830.

E-mail addresses: marco@lce.hut.fi (M. Patriarca), anirban@bnl.gov (A. Chakraborti), kimmo.kaski@hut.fi (K. Kaski).

some simple systems of closed economy, which can be either solved exactly or simulated numerically, in order to investigate the relation between the micro-dynamics and the resulting macroscopic money distribution [7–12]. In this paper we consider the generalization of a simple model of money conserving economy, realized by introducing a criterion of saving in the transaction law, through the saving propensity λ . We study numerically its asymptotic money distribution as a function of the model parameters. We show that it is not a Gibbs distribution and, by direct comparison with numerical data, that the corresponding analytical solution has the form of a γ -distribution.

2. Model

In the simple model considered [8], N agents can exchange money in pairs between themselves. For the sake of simplicity we assume that all the agents are initially assigned the same money amount m_0 , despite this condition is not restrictive for the following results. Agents are then let to interact. At every “time step”, a pair (i, j) is randomly chosen and the transaction takes place. During the transaction, the agent money amounts m_i and m_j undergo a variation, $m_i \rightarrow m'_i$ and $m_j \rightarrow m'_j$. Money is assumed to be conserved during the transaction, so that

$$m_i + m_j = m'_i + m'_j. \quad (1)$$

In this basic model, m'_i and m'_j are obtained through a random reassignment of the total initial money $(m_i + m_j)$,

$$\begin{aligned} m'_i &= \varepsilon(m_i + m_j), \\ m'_j &= (1 - \varepsilon)(m_i + m_j), \end{aligned} \quad (2)$$

where ε is a random number, extracted from a uniform distribution in the interval $(0, 1)$. Note that this model of dynamics, as well as its variations considered in the following, ensures that agents have no debts after the transaction, i.e., they are always left with a money amount $m \geq 0$. It can be shown that, merely as a consequence of the conservation law (1), the system relaxes toward an equilibrium state characterized by a Gibbs distribution [7–9],

$$f(m) = \beta \exp(-\beta m), \quad (3)$$

where $\beta = 1/\langle m \rangle$ represents the inverse average money and $\langle m \rangle = \sum_i m_i/N \equiv m_0$. This means that, after relaxation, the majority of the agents has a very small amount of money, while the number of richest agents—e.g. those with m larger than a given value m' , as well as the fraction of the total money they own, exponentially decreases with m' . The Gibbs distribution (3) has been shown to represent a robust equilibrium state, reached independently of the initial conditions also in generalized models, such as those involving multi-agent transactions.

However, if a saving criterion is introduced [7,9], i.e., agents save a fraction λ —the saving propensity—of the money they have before the transaction is made, the shape of the equilibrium distribution changes dramatically. The conservation equation (1) still




Fig. 1. Equilibrium money distributions for different values of the saving propensity λ , in the closed economy model defined by Eqs. (5). The continuous curves are the fitting functions, defined in Eq. (8). Distributions are normalized to the channel width $dx = 1/N_c = 0.05$, where $N_c = 20$ is the number of channels in the unit interval.

holds, but the money to be shared in a transaction between the i th and the j th agent is now $(1 - \lambda)(m_i + m_j)$. Then Eqs. (2) are thus modified

$$\begin{aligned}
 m'_i &= \lambda m_i + \varepsilon(1 - \lambda)(m_i + m_j), \\
 m'_j &= \lambda m_j + (1 - \varepsilon)(1 - \lambda)(m_i + m_j).
 \end{aligned}
 \tag{4}$$

These equations can also be rewritten in the following way:

$$\begin{aligned}
 m'_i &= m_i + \Delta m, \\
 m'_j &= m_j - \Delta m, \\
 \Delta m &= (1 - \lambda)[\varepsilon m_j - (1 - \varepsilon)m_i],
 \end{aligned}
 \tag{5}$$

which clearly shows how money is conserved during the transaction.

We performed numerical simulations, for various values of λ , of a system with $N = 500$ agents. In each simulation a sufficient number of transactions, as far as 10^7 , depending on the value of λ , was used in order to reach equilibrium. The final equilibrium distributions for a given λ , obtained by averaging over 1000 different runs, are shown in Fig. 1.

3. Fitting

The exact solution for the case $\lambda = 0$ is known to be given by the Gibbs distribution, Eq. (3). Here we give the corresponding exact solution for a generic value of λ , with




Fig. 2. As in Fig. 1, but on a double logarithmic scale.

$0 < \lambda < 1$. This solution was found by fitting the results of numerical simulations and it turns out to fit extremely well all data.

It is convenient to introduce the reduced variable

$$x = \frac{m}{\langle m \rangle}, \tag{6}$$

the agent money in units of the average money $\langle m \rangle$, and the parameter

$$n(\lambda) = 1 + \frac{3\lambda}{1 - \lambda}. \tag{7}$$

We found that the money distributions, for arbitrary values of λ , are well fitted by the function

$$P_n(x) = a_n x^{n-1} \exp(-nx), \tag{8}$$

where x and n are defined in Eqs. (6) and (7), respectively.¹ From the normalization condition, the prefactor is easily shown to be

$$a_n = \frac{n^n}{\Gamma(n)}, \tag{9}$$

where $\Gamma(n)$ is the Gamma function.

The fitting curves for the distribution (continuous lines) are compared with the numerical data in Fig. 1. The fitting describes the distribution also at large values of x , as shown by the logarithmic plots in Fig. 2. The numerical values of the parameters a_n and n are compared with the respective fitting functions (9) and (7) in Fig. 3. The distribution function (8) still contains an exponential factor $\exp(-nx)$, similar to

¹ An excellent fitting is also obtained by replacing $\exp(-nx)$ with $\exp(-nx^c)$, where c is an additional parameter. Since the fitting provides a value c very close to one and is good for all λ 's, we assumed $c \equiv 1$.




Fig. 3. The parameters a_n (left) and n (right) versus λ obtained from numerical data (dots) and the corresponding analytical formulas (continuous curves) given by Eqs. (9) and (7), respectively.

that of the Gibbs distribution, but the average value is now rescaled by n . However, it is the power x^{n-1} which qualitatively changes the shape of the Gibbs distribution, which turns into a curve with a maximum at $x > 0$, i.e., with a mode different from zero.

It is to be noticed that by introducing the rescaled variable

$$x_n = nx \equiv \frac{m}{\langle m \rangle / n} \tag{10}$$

and the corresponding probability density $P_n(x_n) = dF_n(x)/dx_n \equiv P_n(x)/n$, where F_n is the cumulative function, and using the explicit expression of a_n , Eq. (9), distribution (8) becomes

$$P_n(x_n) = \frac{x_n^{n-1} \exp(-x_n)}{\Gamma(n)} \equiv \gamma_n(x_n) \tag{11}$$

which is just the γ -distribution function $\gamma_n(x_n)$ for the variable x_n .

4. Conclusions and discussion

We have studied a generalization of the simple closed economy model, in which a random reassignment of the agent money takes place, by introducing a saving propensity $\lambda > 0$. We suggest a form for the corresponding exact analytical solution, which we have successfully tested through a fitting of the numerical data. The distribution naturally lends itself to be interpreted as a γ -distribution $\gamma_n(x_n)$ for the reduced variable $x_n = m/(\langle m \rangle / n)$. The parameter $n(\lambda) = 1 + 3\lambda/(1 - \lambda)$ is in principle continuous, but it can vary between 1 and ∞ when λ varies between 0 and 1. This result naturally encourages us to provide a deeper physical interpretation and a rigorous derivation of the solution, which are underway.

Acknowledgements

This work was partially supported by the Academy of Finland, Research Center for Computational Science and Engineering, Project no. 44897 (Finnish Centre for Excellence Program 2000–2005). The work at Brookhaven National Laboratory was carried out under Contract no. DE-AC02-98CH10886, Division of Material Science, US Department of Energy.

References

- [1] V. Pareto, *Cours d'economie politique*, Rouge, Lausanne and Paris, 1897.
- [2] M. Levy, S. Solomon, *Physica A* 242 (1997) 90.
- [3] A. Drăgulescu, V.M. Yakovenko, *Physica A* 299 (2001) 213.
- [4] W.J. Reed, B.D. Hughes, *Phys. Rev. E* 66 (2002) 067103.
- [5] H. Aoyama, W. Souma, Y. Fujiwara, *Physica A* 324 (2003) 352.
- [6] F. Slanina, arXiv:cond-mat/0311235, 2003.
- [7] A. Chakraborti, B.K. Chakrabarti, *Eur. Phys. J. B* 17 (2000) 167.
- [8] A. Drăgulescu, V.M. Yakovenko, *Eur. Phys. J. B* 17 (2000) 723.
- [9] A. Chakraborti, *Int. J. Mod. Phys. C* 13 (2002) 1315.
- [10] B. Hayes, *Am. Sci.* 90 (2002) 400.
- [11] A. Chatterjee, B.K. Chakrabarti, S.S. Manna, *Physica A* 335 (2004) 155.
- [12] A. Das, S. Yarlagadda, arXiv:cond-mat/0310343, 2003.